

Suppression of Rat Bone Marrow Cells by Friend Murine Leukemia Virus Envelope Proteins

Stefan Mazgareanu,* Justus G. Müller,† Stefanie Czub,† Simone Schimmer,* Martin Bredt,† and Markus Czub*¹

*Institut für Virologie und Immunbiologie, Universität Würzburg, Versbacherstrasse 7, D-97078 Würzburg; and †Pathologisches Institut, Universität Würzburg, Josef-Schneider-Strasse 2, D-97080 Würzburg, Germany

Received October 1, 1997; returned to author for revision October 30, 1997; accepted December 10, 1997

In a retroviral rat model, we have investigated the nontransforming effects of murine leukemia virus FB29 on the bone marrow. Upon intraperitoneal inoculation with murine leukemia virus FB29 of either neonatal or adult rats, bone marrow cells became massively infected within the first 12 days postinoculation. In neonatally inoculated rats, a persistent productive bone marrow infection was established, whereas in rats inoculated as adults, no infected bone marrow cells could be detected beyond 12 days postinoculation. Retroviral infection was most likely cleared by an antiviral immune response (Hein *et al.*, 1995, *Virology* 211, 408–417). Exposure to virus irreversibly decreased numbers of bone marrow cells staining with monoclonal antibody OX7 by 10–30%. Reduction of OX7⁺ bone marrow cells by 20% was also observed *in vitro*, after bone marrow cells from uninfected adult rats had been co-incubated with virus. FB29-envelope proteins were sufficient alone to reduce numbers of OX7⁺ bone marrow cells, both *in vivo* and *in vitro*. According to results on incorporation of propidium iodide, decreased numbers of OX7⁺ cells were due to cell death. By flow cytometric analyses OX7⁺ bone marrow cells as well as monocytes/macrophages were identified to be major target cells for infection with FB29 within the bone marrow. Thus, the mechanism(s) responsible for death of OX7⁺ bone marrow cells might be due to direct toxicity of viral envelope proteins and/or to interactions of viral envelope proteins with cells of the monocytic lineage. © 1998 Academic Press

INTRODUCTION

Hematological abnormalities are frequently associated with viral infections (Young, 1993). Of the retroviridae, Human Immunodeficiency Virus (HIV) (Re *et al.*, 1994), Simian Immunodeficiency Virus (SIV) (Mandell *et al.*, 1995; Watanabe *et al.*, 1990), and Feline Immunodeficiency Virus (FIV) (Linenberger and Abkowitz, 1995) cause bone marrow disorders with closely resembling characteristics, i.e., peripheral blood cytopenias affecting one or more hematopoietic lineages. This suggests that the pathogenesis of the bone marrow suppression associated with these lentiviruses is similar. Infections with other retroviruses, e.g., Feline Leukemia Virus (Linenberger and Abkowitz, 1995) or Murine Leukemia Virus (MuLV) (Belli and Fan, 1994; Kabat, 1989), often result in proliferative hematological disorders, although other defects originating in the bone marrow have been described (Linenberger and Abkowitz, 1995).

We recently described pathogenic effects of a Friend MuLV, clone FB29, and of a variant, NT40, derived from FB29, in Fisher rats (Czub *et al.*, 1995). Both viruses as well as the related clone PVC-211 (Masuda *et al.*, 1992) cause a progressive neurodegenerative disease in neonatally inoculated Fisher rats. However, neither Friend virus nor the variants induced erythroleukemias in rats,

as they do in mice (Masuda *et al.*, 1992; Sitbon *et al.*, 1986). As many MuLV-induced leukemias in mice depend on infection of one or more hematopoietic lineages, the bone marrow is a prime target for leukemogenic MuLV (Dexter *et al.*, 1977). It was therefore of interest to determine whether in rats MuLV failed to infect the bone marrow.

In our previous study we observed a marked affinity of both viruses, FB29 and NT40, for cells of the monocytic lineage in rats (Czub *et al.*, 1995). However, neither our group nor another group was able to efficiently infect macrophages with MuLV *in vitro* (Wong *et al.*, 1989). For the sheep lentivirus, visna, it has been suggested that infection of macrophage precursors in the bone marrow played an important role for dissemination of virus by means of migration of infected monocytes into various tissues (Gendelman *et al.*, 1985). A similar mechanism could have taken place in rats infected with FB29 or its derivative NT40. Infection of rat macrophage precursors before a postmitotic fixation occurred would provide a suitable explanation for our previous findings of macrophage susceptibility for MuLV (Czub *et al.*, 1995). Therefore, we sought to identify whether rat macrophage precursors were susceptible for infection with MuLV.

Our results revealed that bone marrow cells (BMC) were early and frequent targets for MuLV FB29 after inoculation of both neonatal and adult Fisher rats. Within the bone marrow, infection was clearly detected in Thy 1⁺ cells—among which are early hematopoietic progenitor cells (van den Brink *et al.*, 1991)—as well as in

¹To whom correspondence and reprint requests should be addressed. Fax: +49-(0)931-201 3934. E-mail: czub@vim.uni-wuerzburg.de.




FIG. 1. Course of productive infection with MuLV FB29 in the bone marrow upon intraperitoneal inoculation of neonatal (\diamond) or adult (\blacktriangle) rats. Each symbol represents a titer from an individual animal determined by an infectious center assay.

macrophages. After exposure to MuLV FB29, a Thy 1⁺ subpopulation of rat BMC diminished irreversibly. As FB29-envelope proteins alone were also sufficient to reduce numbers of Thy 1⁺ cells, suppression of Thy 1⁺ BMC was not dependent on the addition of infectious virus, excluding direct infection as an obligatory cause for decreased Thy 1⁺ cell numbers.

RESULTS

Rat BMC are early targets for MuLV FB29

Inoculation of neonatal Fisher rats with MuLV FB29 has been shown to result in persistent infection of several organs such as blood, spleen, thymus, and central nervous system (Czub *et al.*, 1995). Here, we show by means of a kinetic analysis that upon neonatal inoculation of rats BMC became productively infected at around 6–8 days postinoculation (dpi). The number of productively infected BMC increased to levels of 1–10% of all BMC producing virus [$\log 4.8 \pm 0.1$ focus forming units (FFU)/10⁶ BMC; $n = 34$; means \pm SE]. This occurred at around 12–14 dpi and this proportion of infected BMC was lifelong (Fig. 1).

Inoculation of adult Fisher rats with FB29 has previously been shown to result in an effective antiviral humoral and cellular immune response (Hein *et al.*, 1995). In those animals, neither viremia nor productive infection of spleen or thymus has been detected when analyzed at or after 28 dpi (Hein *et al.*, 1995). However, early after inoculation of adult rats, BMC became massively infected and in the first instance, i.e. up to 12 dpi, the kinetics of BMC infection closely resembled that of neonatally inoculated rats (Fig. 1). At no time was viremia detected (data not shown). After 12 dpi, a substantial

reduction of virally infected cells appeared to have occurred, as productively infected cells could not be found either in the bone marrow (Fig. 1) or in other organs of animals inoculated as adults (Hein *et al.*, 1995). It is likely that the decline of virus-infected cells was due to the emerging antiviral immune response. Despite the viral clearance in this latter group of animals, rat BMC were an early as well as a frequent target for MuLV FB29 in both neonatal and adult Fisher rats.

FB29-infected OX7⁺ as well as OX7⁻/CD11b/c⁺ BMC

To identify and characterize BMC interacting with FB29, flow cytometric analyses using cell-specific and virus-specific immunolabeling were performed. In neonatally inoculated rats, around 12–31% ($n = 9$) of all OX7⁺ BMC was found to express viral antigens (Fig. 2A). The number of double positive BMC remained at this level throughout the lifetimes. In rats inoculated as adults, a considerable proportion of OX7⁺ BMC were also positive for antiviral staining (4–12%, $n = 4$; Fig. 2B) when analyzed before a substantial antiviral immune response had developed. Of infected rats inoculated either as neonates or adults, a considerable fraction of cells of the monocytic lineage which were defined by expression of CD11b/c (Robinson *et al.*, 1986) could also be stained with antiviral antibodies (10–45% in neonatally inoculated animals, $n = 7$, Fig. 2D; 4–18% in rats inoculated as adults, $n = 4$, Fig. 2E), indicating viral infection of bone marrow-associated monocytes/macrophages. Viral tropism for monocytes/macrophages of other organs has been observed before in this animal model (Czub *et al.*, 1995). After 14 dpi, BMC expressing viral antigens could not be found any longer in rats inoculated as adults. Altogether, these findings correlated well with the results of the infectious center assays (Fig. 1).

Decreased numbers of OX7⁺ BMC in rats exposed to MuLV FB29

To investigate possible consequences of the massive bone marrow infection a flow cytometric analysis using monoclonal antibody OX7 was performed. This antibody stains many rat BMC including relatively primitive hematopoietic precursor cells (van den Brink *et al.*, 1991). In neonatally inoculated rats we observed a decline of OX7⁺ BMC, paralleling the spread of retroviral infection in this organ (Figs. 1 and 3A). In these persistently infected animals, numbers of OX7⁺ BMC never recovered to levels of uninfected age-matched control animals and remained 22% below normal values (Fig. 3A). Similarly, in rats inoculated as adults, numbers of OX7⁺ BMC fell below normal values at the time when BMC became infected with FB29 and levelled out at 10% below numbers of OX7⁺ BMC from uninfected controls (Fig. 3B). Numbers of OX7⁺ BMC in rats inoculated as adults never reached levels of control animals again although—after




FIG. 2. (A–F) Expression of viral proteins in rat BMC. BMC from rats inoculated as neonates (A,D) or adults (B,E) or from age-matched uninoculated animals (C,F) were stained with monoclonal antibodies OX7 (A,B,C) and OX42 (D,E,F), respectively, and gated. Intracellular staining of viral proteins is depicted on the y axis.

an antiviral immune response had been inaugurated—virus-infected cells could not be detected in these rats (Fig. 1). These results show that the cellular composition of rat bone marrow was significantly modified upon retroviral infection. Furthermore, it appears that the virus-induced modification(s) on the cellular arrangement of the bone marrow was irreversible, as viral clearance

from the bone marrow and from other organs did not lead to restoration of OX7⁺ BMC numbers to normal levels.

Histological analyses

Histological analyses of total bone marrow by means of differential staining procedures revealed equal cellu-




FIG. 3. A–C: Time course of numbers of OX7⁺ rat BMC after inoculation of neonates either with MuLV FB29 (A) or with FB29 envelope proteins (C) and of adult rats with MuLV FB29 (B), respectively. Numbers of OX7⁺ rat BMC were determined by flow cytometry. Each symbol represents the ratio of numbers of OX7⁺ rat BMC from two individual animals: from one animal exposed to viral proteins to numbers of OX7⁺ rat BMC from a second, age-matched noninfected/nonexposed littermate.




FIG. 4. A,B: Time course of numbers of OX7⁺ rat BMC *in vitro*. BMC were incubated with MuLV FB29 (A) or with FB29-envelope proteins (B). Numbers of OX7⁺ cells were determined by flow cytometry. Each symbol represents the ratio of numbers of OX7⁺ rat BMC exposed to viral proteins to numbers of OX7⁺ mock-treated cells (shown is a mean of nine experiments).

larity, normal architecture, and maturation of the erythroid, myeloid, and thromboid lineage in infected bone marrow compared to that of normal animals (data not shown). There was no increase in either storage iron or fibrosis in infected bone marrow (data not shown). This indicated that the actual process of Thy 1⁺ cell number reduction was difficult to visualize by histopathological techniques and that apparent reactions of the host in response to the decline of OX7⁺ BMC did not occur.

Rapid decline of OX7⁺ BMC after exposure to MuLV FB29 *in vitro*

To further explore virus-induced bone marrow alterations, BMC from uninfected adult rats (>5 weeks) were incubated *in vitro* with FB29 (MOI 1) or with Mock-in-

fect cell culture supernatant and analyzed with regard to the numbers of OX7⁺ BMC at various timepoints thereafter. Within 48 h of cultivation, numbers of OX7⁺ BMC co-incubated with virus declined by 22% compared to those of control BMC (Fig. 4A). Thus, *in vitro* cultivated BMC responded to virus exposure in a manner similar to BMC *in vivo*, indicating that the virus-induced suppression of OX7⁺ BMC was an intrinsic feature of the bone marrow and independent from, e.g., cellular influx from other organs. Surprisingly, cocultivation of BMC with FB29 resulted in only very limited productive infection of BMC, i.e., $\log 2.4 \pm 0.2$ FFU/10⁶ BMC ($n = 5$; means \pm SE).

FB29 envelope protein(s) induced reduction of OX7⁺ BMC both *in vivo* and *in vitro*

As BMC hardly became infected *in vitro*, however, numbers of OX7⁺ BMC decreased upon viral exposure, and we studied the effects of viral proteins alone. Previous investigations on the pathogenesis of MuLV-induced diseases have pointed to viral envelope protein(s) playing a major role in the disease processes (Hasenkrug *et al.*, 1996; Portis *et al.*, 1990; Sitbon *et al.*, 1986). We constructed a FB29-envelope protein expression vector (Fig. 5A) to test whether viral surface protein(s) alone would be sufficient to reduce numbers of OX7⁺ BMC. Expression and secretion of FB29-envelope proteins into supernatants of transfected 3T3 cells was controlled by flowcytometric (Fig. 5B) and immunoblot (Fig. 5C) analyses. Supernatants of transfected cells containing FB29 envelope proteins were administered intraperitoneally (ip) into neonatal rats. Animals treated with envelope proteins exhibited a reduced number of OX7⁺ BMC com-




FIG. 5. (A–C) Schematic diagram of a FB29-envelope gene expression vector, LTR-env-LTR(pBK) (A), and control of expression of FB29-envelope proteins in 3T3 cells. Expression of FB29-envelope proteins on NIH3T3 cells transfected with LTR-env-LTR(pBK) was analyzed by flow cytometry. Transfected (plain line) and nontransfected (dotted line) cells were stained with mAb 48 specific for gp70 of FB29 (B). Secretion of FB29-envelope proteins from NIH3T3 cells either transfected with LTR-env-LTR(pBK) (*env*) or with a control vector LTR-GFP-LTR(pBK) (*GFP*) and of supernatants from nontransfected cells was analyzed by immunoblotting (C).




FIG. 6. (A,B) Time course of numbers of OX7⁺ (A) and of OX22⁺ (B) rat BMC after inoculation of neonates with MuLV FB29. Numbers of OX7⁺ and of OX22⁺ rat BMC were determined by flow cytometry. Each symbol represents the ratio of numbers of OX7⁺ rat BMC from two individual animals: from one infected animal to numbers from a second, age-matched noninfected littermate. Of OX7⁺ BMC 95% were also positive for mAb OX22 (not shown).

pared to numbers of BMC from animals treated with control supernatants from untransfected cells (Fig. 3C). The decline of OX7⁺ BMC after treatment with envelope proteins was not as strong as after inoculation of replication-competent virus into neonates; however, just like in infected rats, numbers of OX7⁺ BMC from rats treated with envelope proteins remained below numbers of OX7⁺ BMC from control animals for life. Infectious virus bearing an envelope protein reactive with mAb 48 was never recovered from rats inoculated with viral proteins (data not shown), excluding the possibilities of laboratory contamination or generation of new viruses. In *in vitro* experiments, FB29-envelope proteins also reduced numbers of OX7⁺ BMC by up to 24% (Fig. 4B). These results demonstrate that infection was not obligatory for the induction of bone marrow alterations and that viral envelope proteins were sufficient to suppress numbers of OX7⁺ BMC.

Decline of OX7⁺ BMC was due to cell death

If the decline of OX7⁺ BMC was due to downregulation of Thy 1 molecules from the cell surface, other cellular surface markers on these cells might not be affected after exposure to viral proteins. Therefore, we doubly labeled BMC with mAb OX7 and with mAb OX22, the latter staining 95% of OX7⁺ BMC (data not shown). In neonatally inoculated animals, numbers of OX7⁺ and of OX22⁺ BMC declined in a similar manner, when stained in parallel at various timepoints pi (Figs. 6A and 6B). This suggested a loss of cells rather than a loss of cellular surface marker Thy 1.

To determine whether OX7⁺ BMC died upon exposure to FB29-envelope proteins we analyzed BMC in our *in vitro* system by means of propidium iodide (PI) incorporation. Thirty hours post initial exposure, OX7⁺ BMC exposed to viral envelope proteins incorporated PI more frequently than OX7⁺ BMC treated with control supernatants (Fig. 7). This tendency increased

with time (Fig. 7). These results demonstrate that retroviral envelope proteins exhibited cytotoxic effects on Thy 1⁺ BMC from rats.

DISCUSSION

Many retroviruses can cause hematopoietic abnormalities, including HIV in humans (Maciejewski *et al.*, 1994; Re *et al.*, 1994), SIV in monkeys (Mandell *et al.*, 1995; Watanabe *et al.*, 1990), FIV in cats (Linenberger and Abkowitz, 1995), and MuLV in mice (Li and Fan, 1990; Tse *et al.*, 1994). Here we report that retroviral infection of rats also caused bone marrow alterations. In all of our experimental approaches using MuLV FB29, we observed a reduction of numbers of Thy 1⁺ BMC, following very closely viral replication and/or the presence of viral envelope proteins.

Both Thy 1⁺ cells and monocytes/macrophages were identified as major cellular targets within the bone marrow for FB29. The Thy 1 antigen, a glycoprotein of 25 kDa (Letarte-Muirhead *et al.*, 1975), is present on 25–40% of rat BMC (Williams, 1976). Cells bearing this antigen represent immature hematopoietic cells, including pluripotent stem cells (van den Brink *et al.*, 1991). Both infection and decline of Thy 1⁺ BMC targeted only fractions of these cells but not the entire population, indicating that a subpopulation(s) with high susceptibility for infection and/or alteration existed. Once Thy 1⁺ subpopulations receptive for infection, alteration, or both are defined more precisely, it will be possible to show whether the decline of Thy 1⁺ BMC is due to direct interaction with viral envelope proteins or to indirect mechanisms.

In addition to cells bearing the Thy 1 antigen, mono-




FIG. 7. Effects of FB29-envelope proteins on the viability of Thy 1⁺ BMC *in vitro*. BMC were incubated with FB29-envelope proteins (▲) or with tissue culture supernatants from NIH3T3 cells transfected with LTR-GFP-LTR (◇). Thy 1⁺ BMC were analyzed for their capacity to incorporate propidium iodide (PI) at different timepoints. Each symbol represents the ratio of Thy 1⁺/PI⁺ BMC to Thy 1⁺/PI⁻ BMC. With time, Thy 1⁺ BMC exposed to viral envelope proteins incorporated PI more frequently than OX7⁺ BMC treated with control supernatants. Shown is a representative result of a total of three experiments.

cytes/macrophages were found to be susceptible for infection with FB29 within the bone marrow. Infection of bone marrow resident monocytes/macrophages went along without apparent modification of these cells. We did not observe either macrophage hyperplasia or myeloid hyperplasia as described in bone marrows of monkeys early after SIV infection (Mandell *et al.*, 1995). However, cells of the monocytic lineage often become activated upon retroviral infection, with the possible consequence of damaging neighboring cells. For example, HIV-induced bone marrow failure appears to be related to TNF- α secretion by macrophages (Maciejewski *et al.*, 1994). Just like after infection with FB29, macrophages are a major cellular target for HIV (McElrath *et al.*, 1989) as well as for SIV (Mandell *et al.*, 1995). Enhanced TNF- α production occurs upon exposure of macrophages to HIV envelope proteins gp120/160 (Clouse *et al.*, 1991; Maciejewski *et al.*, 1994; Riekmann *et al.*, 1991). Effects of enhanced TNF- α levels on BMC could be of an inhibitory (Jacobsen *et al.*, 1994) or a cytotoxic nature (Bigda *et al.*, 1994) or both. Similarly, numbers of Thy 1⁺ BMC might have been reduced by an indirect mode of actions mediated through monocytes/macrophages activated by FB29-envelope proteins.

We would also expect pathological long-term consequences of high levels of TNF- α . As such, differentiation of the bone marrow could shift toward monocytes and macrophages, as observed *in vitro* (Jacobsen *et al.*, 1994; Murphy *et al.*, 1988). It would be interesting to find a retroviral-induced TNF- α -mediated shift of hematopoietic differentiation toward monocytes and macrophages *in vivo*, too, because macrophagetropic viral variants might take advantage of such a situation for enhanced viral spread and subsequent further tissue damage, including the bone marrow, in later stages of infection. This hypothesis could also explain why we observed an even more dramatic decline of Thy 1⁺ BMC beginning at 40–50 days after FB29 infection or administration of FB29-envelope proteins into neonatal animals.

Both *in vivo* and *in vitro* data revealed that FB29-envelope proteins alone were sufficient to reduce numbers of Thy 1⁺ BMC. Analyses based on PI incorporation demonstrated that cell death was responsible for the decline of numbers of OX7⁺ BMC. It is thus likely that FB29-envelope proteins are directly cytotoxic. Apart from bone marrow suppression, cytotoxic effects of retroviral envelope proteins might also be involved in other pathological alterations, such as hemolytic anemia of mice (Sitbon *et al.*, 1986), neurodegeneration of both humans and rodents (Hasenkrug *et al.*, 1996; Lipton *et al.*, 1991; Portis *et al.*, 1990; Toggas *et al.*, 1994), and T-cell loss during AIDS (Siliciano, 1996).

It has been suggested that antibodies directed against HIV-envelope proteins are capable of inhibiting colony formation by bone marrow progenitor cells derived from AIDS patients (Donahue *et al.*, 1987). An immune-medi-

ated mechanism might also play an important role in Parvovirus-induced myeloid depression (Segovia *et al.*, 1995). We found no additional suppressive effects on numbers of Thy 1⁺ BMC after clearance of viral infection from rats inoculated as adults. This indicates that elimination of virus-infected BMC by an antiviral immune response was compensated well in these animals. Furthermore, as the reduction of Thy 1⁺ BMC was irreversible, clearance of viral infection did not have a beneficial influence on the numbers of Thy 1⁺ BMC. Therefore, long-term consequences of reduced numbers of Thy 1⁺ BMC might be equal in all animals exposed to virus or viral envelope proteins—independently of their state of infection and the antiviral immune response.

In summary, we describe a new bone marrow abnormality induced by envelope proteins of Friend MuLV in rats. As in susceptible mice, the rat bone marrow was shown to be a major target organ for MuLV; however, it does not give rise to leukemias. We further demonstrated lifelong susceptibility of the rat bone marrow for at least transient FB29 infection and associated alterations. This contrasted with the situation for MuLV-induced leukemogenesis in AKR mice (Buckheit *et al.*, 1987, 1988) and in mice inoculated with Moloney MuLV (Belli and Fan, 1994), respectively. In both cases, early postnatal infection of the bone marrow with MuLV has been proposed to be crucial for the development of leukemia in mice. It also appeared that the reduction of numbers of Thy 1⁺ BMC in rats very closely followed viral infection and/or exposure to viral proteins but that the pathogenesis of MuLV-induced leukemia in mice is composed of a rather slow multistep process (Belli and Fan, 1994; Kabat, 1989; Li and Fan, 1990). Thus, in the course of MuLV-induced rat bone suppression quite different virus–host interactions appear to be recruited than in the pathogenesis of MuLV-induced leukemias in mice.

MATERIALS AND METHODS

Animals, inoculation, viruses, and infectious center assay

Fisher rats were initially obtained from the Institut für Versuchstierzucht (Hannover, Germany) and were bred at our institute. Neonatal rats (0–24 hours postpartum) were inoculated *ip* with 100- μ l virus stock (containing 5.0×10^4 – 10^5 FFU; adult rats (>5 weeks) were inoculated *ip* with 500- μ l virus stock. For some experiments, neonatal rats were inoculated with 100 μ l of NIH3T3-cell culture supernatant transfected with FB29-envelope gene expression vector LTR-env-LTR(pBK) and with LTR-GFP-LTR(pBK), respectively.

FB29 is a molecular clone of the I⁻⁵ strain of Friend MuLV (Sitbon *et al.*, 1986). Neonatal inoculation of rats with FB29 leads to infection of the brain, resulting in spongiform neurodegeneration and mild neurological signs (Hein *et al.*, 1995). Infectivity of cell culture and of

BMC was determined by means of a focal immunoassay using monoclonal antibody (mAb) 48 specific for FB29 (Robertson *et al.*, 1991), peroxidase-conjugated rabbit anti-mouse immunoglobulin (Dako), and 3-amino-9-ethyl-carbazol (Sigma) as described previously (Czub *et al.*, 1991).

Construction of a FB29-envelope gene expression vector, LTR-env-LTR(pBK)

An envelope-LTR fragment (*XbaI*–*PstI*) from molecular clone FB29 (Genbank Accession No. Z11128) encompassing nucleotides 5634–561 was cloned into pBKCMV (Stratagene) from which the CMV promoter (*SpeI*–*NheI*) had been excised before. A 5' LTR (*ClaI*–*PstI*) encompassing nucleotides 7702–561 from FB29 was placed upstream of the envelope-LTR sequence in pBK. A neomycin resistance gene outside retroviral vector sequences is driven by a SV40 promoter (Fig. 5A). A similar procedure was chosen for construction of control vector LTR-GFP-LTR(pBK) in which the envelope gene was replaced by the green fluorescent protein, GFP (Heim *et al.*, 1995). Plasmid DNA was transfected into NIH3T3 cells using the calcium phosphate procedure (Graham and van der Eb, 1973). Cells were cultivated in 10% FCS–RPMI containing 800 $\mu\text{g/ml}$ G418 (Gibco-BRL) to select for stably transfected cell clones. Supernatants were taken 2 weeks after withdrawal of G418 from the medium.

Cultivation of BMC

BMC were obtained by flushing the marrow cavity with phosphate-buffered saline (PBS). The cells were washed, lysed with lysis buffer (0.155 M NH_4Cl , 0.001 M KH_2CO_3 , 0.1 mM EDTA in H_2O) for 4 min on ice, washed again, and counted. A total of 5.0×10^5 BMC per well of a TC12-plate (Nunc, Denmark) was seeded in 1.0 ml 10% FCS–RPMI 1640 containing 72 mg/l penicillin, 120 mg/l streptomycin, and 0.0028 M glutamine. Cells were incubated for 24 h at 37°C in a humidified atmosphere containing 5 vol.-% CO_2 . After 24 h, MuLV stock or cell culture supernatant from either MOCK infected or transfected cells was added to BMC cultures. At 12, 24, 36, and 48 h after initial exposure, BMC were resuspended by intensive pipetting and immunostained with mAb OX7 specific for Thy 1 (Mason and Williams, 1980) and phycoerythrin-conjugated goat anti-mouse immunoglobulin-F(ab)-fragment (Dianova) as described previously (Hein *et al.*, 1995).

Flow cytometry analysis

NIH3T3 cells stably transfected with LTR-env-LTR(pBK) or BMC were analyzed with a FACScan (Becton-Dickinson). A total of 2.0×10^5 cells were washed in FACS buffer (0.1% bovine serum albumin + 0.02% NaN_3 in PBS). BMC were stained with biotinylated mAb OX7 (Di-

anova), Texas Red 670-conjugated streptavidin (Gibco-BRL), and with fluorescein isothiocyanate(FITC)-conjugated mAb OX42 specific for CD11b/c (Robinson *et al.*, 1986) or with mAb OX22 reacting with the high-molecular-weight forms of CD45 (Spickett *et al.*, 1983). NIH3T3 cells stably transfected with LTR-env-LTR(pBK) were analyzed with mAb 48 and phycoerythrin-conjugated goat anti-mouse immunoglobulin G-F(ab)-fragment (Dianova). Intracellular staining for virus proteins in BMC was performed after staining of the cell surface. Cells were fixed with formaldehyde (3.7%) for 30 min at room temperature, permeabilized with Triton X-100 (0.3%) (Sigma) for 30 min at room temperature, and stained intracellularly with goat polyclonal antiserum directed against MuLV gp70 (kindly donated by R. Friedrich, Gießen) and FITC-conjugated rabbit anti-goat immunoglobulin G (Dako). Control staining was performed with normal goat serum.

Propidium iodide staining

In some experiments, BMC were incubated with PI (20 $\mu\text{g/ml}$) for 10 min at 4°C and analyzed by flow cytometry to distinguish dead from live cells (Radosovic *et al.*, 1990). Prior to PI incubation cells were labeled with mAb OX7.

Immunoblot analysis

NIH3T3 cells were lysed in lysis buffer [4% sodium-dodecyl-sulfate, 20% glycerin, 16.6 mM Tris (pH 6.8), and 4% β -mercaptoethanol] and boiled for 10 min. Supernatants of transfected or nontransfected cells were precipitated with trichloroacetic acid (final concentration 5%, 10 min on ice), pelleted (10 min, 5000g), washed with methanol, dried, resuspended in equal volumes of PBS and 2 \times lysis buffer, and boiled for 10 min. A total of 10 μl of the samples were subjected to electrophoresis in SDS-polyacrylamide gel and subsequently transferred onto a nitrocellulose filter. Nitrocellulose filters were blocked for 1 h at room temperature with PBS–Tween (0.1% bovine serum albumin, 3% nonfat powdered milk, 0.5% Tween 20) and incubated overnight at 4°C with goat polyclonal antiserum directed against MuLV gp70 (1/3000) and peroxidase-coupled rabbit anti-goat immunoglobulin G antibodies (Dianova) (1/1000). Nitrocellulose filters were washed twice in PBS–Tween for 10 min at room temperature before revelation with the chemiluminescence ECL kit (Amersham Corp.) following the recommendations of the manufacturer.

Histological examinations

Sections of 5 μm bone marrow were stained with H&E, Giemsa, DiPAS, the Prussian blue iron stain, and the Gomory silver staining method. By light microscopy, the following parameters were evaluated: overall cellularity, maturation in the myelopoiesis, erythropoiesis, thrombopoiesis, the medullary architecture, and the reticulin fiber

content. The evaluation of the bone included the architecture of the spongiosa and the presence of osteoblasts and osteoclasts on the surface.

ACKNOWLEDGMENTS

We thank A. Kolb and J. Herold for help with the construction of the retroviral vectors, E. Bachmann for excellent technical assistance, and C. Jassoy, S. Niewiesk, and V. ter Meulen for critical review of the manuscript. This work was supported by grants from the Deutsche Forschungsgemeinschaft (CZ 56/1-2) and from the Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie (II-068-88), Germany.

REFERENCES

- Belli, B., and Fan, H. (1994). The leukemogenic potential of an enhancer variant of moloney murine leukemia virus varies with the route of inoculation. *J. Virol.* **68**, 6883–6889.
- Bigda, J., Beletsky, I., Brakebush, C., Varfolomeev, Y., Engelmann, H., Bigda, J., Holtmann, H., and Wallach, D. (1994). Dual role of the p75 tumor necrosis factor (TNF) receptor in TNF cytotoxicity. *J. Exp. Med.* **180**, 445–460.
- Buckheit, R. W., Bolognesi, D. P., and Weinhold, K. J. (1987). The effects of leukemiosuppressive immunotherapy on bone marrow infectious cell centers in AKR mice. *Virology* **157**, 387–396.
- Buckheit, R. W., Kurtzberg, J., Bolognesi, D. P., and Weinhold, K. J. (1988). The coenrichment of stem cells, prothymocytes, and stromal elements with ecotropic retrovirus-producing cells from the bone marrow of leukemia-prone AKR mice. *Virology* **162**, 354–361.
- Clouse, K. A., Cosentino, M. L., Weih, K. A., Pyle, S. W., Robbins, P. B., Hochstein, H. D., Natarjan, V., and Farrar, W. (1991). The HIV-1 gp120 envelope protein has the intrinsic capacity to stimulate monokine secretion. *J. Immunol.* **147**, 2892–2901.
- Czub, M., Czub, S., McAtee, F. J., and Portis, J. L. (1991). Age-dependent resistance to murine retrovirus-induced spongiform neurodegeneration results from central nervous system-specific restriction of virus replication. *J. Virol.* **65**, 2539–2544.
- Czub, M., Czub, S., Rappold, M., Mazgareanu, S., Schwender, S., Demuth, M., Hein, A., and Dörries, R. (1995). Murine leukemia virus induced neurodegeneration of rats: Enhancement of neuropathogenicity correlates with enhanced viral tropism for macrophages, microglia, and brain vascular cells. *Virology* **214**, 239–244.
- Dexter, T. M., Scott, D., and Teich, N. M. (1977). Infection of bone marrow cells in vitro with FLV: Effects on stem cell proliferation, differentiation and leukemogenic capacity. *Cell* **12**, 355–364.
- Donahue, R. E., Johnson, M. M., Zon, L. I., Clark, S. C., and Groopman, J. E. (1987). Suppression of *in vitro* haematopoiesis following human immunodeficiency virus infection. *Nature* **326**, 200–203.
- Gendelman, H. E., Narayan, O., Molineaux, S., Clements, J. E., and Ghotbi, Z. (1985). Slow, persistent replication of lentiviruses: Role of tissue macrophages and macrophage precursors in bone marrow. *Proc. Natl. Acad. Sci. USA* **82**, 7086–7090.
- Graham, F. L., and van der Eb, A. J. (1973). A new technique for the assay of infectivity of human adenovirus 5 DNA. *Virology* **52**, 456–467.
- Hasenkrug, K. J., Robertson, S. J., Portis, J. L., McAtee, F., Nishio, J., and Chesebro, B. (1996). Two separate envelope regions influence induction of brain disease by a polytropic murine retrovirus (FMCF98). *J. Virol.* **70**, 4825–4828.
- Heim, R., Cubitt, A. B., and Tsien, R. Y. (1995). Improved green fluorescence. *Nature* **373**, 664–665.
- Hein, A., Czub, S., Xiao, L. X., Schwender, S., Dörries, R., and Czub, M. (1995). Effects of adoptive immune transfers on murine leukemia virus-infection of rats. *Virology* **211**, 408–417.
- Jacobsen, F. W., Rothe, M., Rusten, L., Goeddel, D. V., Smeland, E. B., Veiby, O. P., Stordal, L., and Jacobsen, S. E. W. (1994). Role of the 75-kDa tumor necrosis factor receptor: Inhibition of early hematopoiesis. *Proc. Natl. Acad. Sci. USA* **91**, 10695–10699.
- Kabat, D. (1989). Molecular biology of friend viral erythroleukemia. *CTMI* **148**, 1–42.
- Letarte-Muirhead, M., Barclay, A. N., and Williams, A. F. (1975). Purification of the Thy-1 molecule, a major cell-surface glycoprotein of rat thymocytes. *Biochem. J.* **151**, 685–697.
- Li, Q., and Fan, H. (1990). Combined infection by moloney murine leukemia virus and a mink cell focus-forming virus recombinant induces cytopathic effects in fibroblasts or in long-term bone marrow cultures from preleukemic mice. *J. Virol.* **64**, 3701–3711.
- Linenberger, M. L., and Abkowitz, J. L. (1995). Haematological disorders associated with feline retrovirus infections. *Baillieres Clin. Haematol.* **8**, 73–112.
- Lipton, S. A., Sucher, N. J., Kaiser, P. K., and Dreyer, E. B. (1991). Synergistic effects of HIV coat protein and NMDA receptor-mediated neurotoxicity. *Neuron* **7**, 11–118.
- Maciejewski, J. P., Weichold, F. F., and Young, N. S. (1994). HIV-1 suppression of hematopoiesis in vitro mediated by envelope glycoprotein and TNF- α . *J. Immunol.* **153**, 4303–4310.
- Mandell, C. P., Jain, N. C., Miller, C. J., and Dandekar, S. (1995). Bone marrow monocytes/macrophages are an early cellular target of pathogenic and nonpathogenic isolates of simian immunodeficiency virus (SIVmac) in rhesus macaques. *Lab. Invest.* **72**, 323–333.
- Mason, D. W., and Williams, A. F. (1980). The kinetics of antibody binding to membrane antigens in solution and at the cell surface. *Biochem. J.* **187**, 1–20.
- Masuda, M., Remington, M. P., Hoffman, P. M., and Ruscetti, S. K. (1992). Molecular characterization of a neuropathogenic and nonerythro-leukemogenic variant of friend murine leukemia virus PVC-211. *J. Virol.* **66**, 2798–2806.
- McElrath, M. J., Pruett, J. E., and Cohn, Z. A. (1989). Mononuclear phagocytes of blood and bone marrow: Comparative roles as viral reservoirs in human immunodeficiency virus type 1 infections. *Proc. Natl. Acad. Sci. USA* **86**, 675–679.
- Murphy, M., Perussia, B., and Trinchieri, G. (1988). Effects of recombinant tumor necrosis factor, lymphotoxin, and immune interferon on proliferation and differentiation of enriched hematopoietic precursor cells. *Exp. Hematol.* **16**, 131–138.
- Portis, J. L., Czub, S., Garon, C. F., and McAtee, F. J. (1990). Neurodegenerative disease induced by the wild mouse ecotropic retrovirus is markedly accelerated by long terminal repeat and *gag-pol* sequences from nondefective friend murine leukemia virus. *J. Virol.* **64**, 1648–1656.
- Radosevic, K., Garritsen, H. S. P., van Graft, M., de Grooth, B. G., and Greve, J. (1990). A simple and sensitive flow cytometric assay for the determination of the cytotoxic activity of human natural killer cells. *J. Immunol. Methods* **135**, 81–89.
- Re, M. C., Furlini, G., Zauli, G., and La Placa, M. (1994). Human immunodeficiency virus type 1 (HIV-1) and human hematopoietic progenitor cells. *Arch. Virol.* **137**, 1–23.
- Riekman, P., Poli, G., Fox, H. C., Kehrl, J. H., and Fauci, A. S. (1991). Recombinant gp120 specifically enhances tumor necrosis factor production and IgG secretion from B lymphocytes from HIV-infected individuals but not from seronegative donors. *J. Immunol.* **147**, 2992–2927.
- Robertson, M. N., Miyazawa, M., Mori, S., Caughey, B., Evans, L., Hayes, S. F., and Chesebro, B. (1991). Production of monoclonal antibodies reactive with a denatured form of the Friend murine leukemia virus gp70 envelope protein: Use in a focal infectivity assay, immunohistochemical studies, electron microscopy and Western blotting. *J. Virol. Methods* **34**, 255–271.
- Robinson, A. P., White, T. M., and Mason, D. W. (1986). Macrophage heterogeneity in the rat as delineated by two monoclonal antibodies MRC OX-41 and MRC OX-42, the latter recognizing complement receptor type 3. *Immunology* **57**, 239–247.

- Segovia, J. C., Bueren, J. A., and Almendral, J. M. (1995). Myeloid depression follows infection of susceptible newborn mice with the parvovirus minute virus of mice (Strain i). *J. Virol.* **69**, 3229–3232.
- Siliciano, R. F. (1996). The role of CD4 in HIV envelope-mediated pathogenesis. *CTMI* **206**, 159–179.
- Sitbon, M., Sola, B., Evans, L., Nishio, J., Hayes, S. F., Nathanson, K., Garon, C. F., and Chesebro, B. (1986). Hemolytic anemia and erythrocytopenia, two distinct pathogenic effects of Friend MuLV: mapping of the effects to different regions of the viral genome. *Cell* **47**, 851–859.
- Spickett, G. P., Brandon, M. R., Mason, D. W., Williams, A. F., and Woollett, G. R. (1983). MRC OX-22, a monoclonal antibody that labels a new subset of T lymphocytes and reacts with the high molecular weight form of the leucocyte-common antigen. *J. Exp. Med.* **158**, 795–810.
- Toggas, S. M., Masliah, E., Rockenstein, E. M., Rall, G. F., Abraham, C. R., and Mucke, L. (1994). Central nervous system damage produced by expression of the HIV-1 coat protein gp120 in transgenic mice. *Nature* **367**, 188–193.
- Tse, K.-F., Morrow, J. K., Hughes, N. K., and Gallicchio, V. S. (1994). Stromal cell lines derived from LP-BM5 murine leukemia virus-infected long-term bone marrow cultures impair hematopoiesis in vitro. *Blood* **84**, 1508–1518.
- van den Brink, M. R. M., Herberman, R. B., and Hiserodt, J. C. (1991). Generation of natural killer cells from Thy 1.1+ bone marrow precursor cells in the rat. *Blood* **2392–2395**.
- Watanabe, M., Ringler, D. J., Nakamura, M., DeLong, P. A., and Letvin, N. L. (1990). Simian immunodeficiency virus inhibits bone marrow hematopoietic progenitor cell growth. *J. Virol.* **64**, 656–663.
- Williams, A. F. (1976). Many cells in the rat bone marrow have cell-surface Thy-1 antigen. *Eur. J. Immunol.* **6**, 526–528.
- Wong, P. K. Y., Prasad, G., Hansen, J., and Yuen, P. H. (1989). *ts1*, a mutant of Moloney murine leukemia virus-TB, causes both immunodeficiency and neurologic disorders in BALB/c mice. *Virology* **170**, 450–459.
- Young, N. S., Ed. (1993) "Viruses and Bone Marrow," Vol. 16, "Hematology." Dekker, New York.